

#14335 Store at +4°C

NFAT1 (D43B1) XP® Rabbit mAb (PE Conjugate)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #Q13469	Entrez-Gene Id: 4773
-------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. <i>Do not aliquot the antibodies. Protect from light. Do not freeze.</i>	
Specificity / Sensitivity	NFAT1 (D43B1) XP® Rabbit mAb (PE Conjugate) recognizes endogenous levels of total NFAT1 protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly87 of human NFAT1 protein.	
Product Description	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated NFAT1 (D43B1) XP® Rabbit mAb #5861.	
Background	The NFAT (nuclear factor of activated T cells) family of proteins consists of NFAT1 (NFATc2 or NFATp), NFAT2 (NFATc1 or NFATc), NFAT3 (NFATc4), and NFAT4 (NFATc3 or NFATx). All members of this family are transcription factors with a Rel homology domain and regulate gene transcription in concert with AP-1 (Jun/Fos) to orchestrate an effective immune response (1,2). NFAT proteins are predominantly expressed in cells of the immune system, but are also expressed in skeletal muscle, keratinocytes, and adipocytes, regulating cell differentiation programs in these cells (3). In resting cells, NFAT proteins are heavily phosphorylated and localized in the cytoplasm. Increased intracellular calcium concentrations activate the calcium/calmodulin-dependent serine phosphatase calcineurin, which dephosphorylates NFAT proteins, resulting in their subsequent translocation to the nucleus (2). Termination of NFAT signaling occurs upon declining calcium concentrations and phosphorylation of NFAT by kinases such as GSK-3 or CK1 (3,4). Cyclosporin A and FK506 are immunosuppressive drugs that inhibit calcineurin and thus retain NFAT proteins in the cytoplasm (5).	
Background References	<ol style="list-style-type: none"> 1. Northrop, J.P. et al. (1993) <i>J Biol Chem</i> 268, 2917-23. 2. Hogan, P.G. et al. (2003) <i>Genes Dev</i> 17, 2205-32. 3. Crabtree, G.R. and Olson, E.N. (2002) <i>Cell</i> 109 Suppl, S67-79. 4. Okamura, H. et al. (2004) <i>Mol Cell Biol</i> 24, 4184-95. 5. Shaw, K.T. et al. (1995) <i>Proc Natl Acad Sci U S A</i> 92, 11205-9. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)	
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.	
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in	

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.