#15002 Store at +4C

AML1 (D33G6) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: Reactivity: Sensitivity: Source/Isotype: **UniProt ID:** Entrez-Gene Id: FC-FP H Mk Endogenous Rabbit IgG #Q01196 861 **Product Usage Application** Dilution Information 1:50 Flow Cytometry (Fixed/Permeabilized) Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the **Storage** antibodies. Protect from light. Do not freeze. AML1 (D33G6) XP® Rabbit mAb (PE Conjugate) recognizes endogenous levels of total AML1 protein. Specificity / Sensitivity Source / Purification Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to amino acids at the amino terminus of human AML1 protein. This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct **Product Description** flow cytometry analysis in human cells. The antibody is expected to exhibit the same species crossreactivity as the unconjugated AML1 (D33G6) XP® Rabbit mAb #4336. AML1 (also known as Runx1, CBFA2, and PEBP2αB) is a member of the core binding factor (CBF) family **Background** of transcription factors (1,2). It is required for normal development of all hematopoietic lineages (3-5). AML1 forms a heterodimeric DNA binding complex with its partner protein CBFβ and regulates the expression of cellular genes by binding to promoter and enhancer elements. AML1 is commonly translocated in hematopoietic cancers: chromosomal translocations include t(8:21) AML1-ETO, t(12:21) TEL-AML, and t(8;21) AML-M2 (6). Phosphorylation of AML1 on several potential serine and threonine sites, including Ser249, is thought to occur in an Erk-dependent manner (7,8). 1. Wang, S. et al. (1993) Mol Cell Biol 13, 3324-39. **Background References** 2. Ogawa, E. et al. (1993) Proc Natl Acad Sci U S A 90, 6859-63. 3. Okuda, T. et al. (1996) Cell 84, 321-30. 4. Wang, Q. et al. (1996) Proc Natl Acad Sci U S A 93, 3444-9. 5. North, T.E. et al. (2004) Stem Cells 22, 158-68. 6. Blyth, K. et al. (2005) Nat Rev Cancer 5, 376-87. 7. Tanaka, T. et al. (1996) Mol Cell Biol 16, 3967-79. 8. Zhang, Y. et al. (2004) J Biol Chem 279, 53116-25.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.