

#81464 Store at -20°C

Cyclin T1 (D1B6G) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, ChIP, C&R	H M R Mk	Endogenous	81	Rabbit IgG	#O60563	904

Product Usage Information

For optimal ChIP results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:100
Chromatin IP	1:50
CUT&RUN	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Cyclin T1 (D1B6G) Rabbit mAb recognizes endogenous levels of total cyclin T1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly642 of human cyclin T1 protein.

Background

Positive transcription elongation factor (P-TEFb) is a heterodimer composed of cyclin T proteins and CDK9. P-TEFb plays a critical role in the transition of the RNA polymerase II (RNAPII) machinery from transcription initiation to elongation (1). At some genes during transcription initiation, RNAPII moves approximately 50 nucleotides away from the transcription start site into the gene where it then pauses and awaits signaling for the formation of a productive transcription elongation complex (1,2). The release of this promoter proximal pausing of RNAPII is signaled by phosphorylation of the C-terminal domain (CTD) within the largest subunit of RNAPII at Ser2 of the heptapeptide repeat sequence by P-TEFb (3). This phosphorylation event is important for the recruitment of mRNA processing factors and chromatin modifiers that are necessary for proper gene expression (4,5). P-TEFb also promotes transcription elongation by phosphorylating DSIF (DRB-induced stimulating factor) and NELF (negative elongation factor), two negative elongation factors that retain RNAPII at the promoter proximal region of genes to initiate transcription elongation (6,7).

Background References

1. Cho, S. et al. (2010) *Cell Cycle* 9, 1697-705.
2. Saunders, A. et al. (2006) *Nat Rev Mol Cell Biol* 7, 557-67.
3. Buratowski, S. (2009) *Mol Cell* 36, 541-6.
4. Lenasi, T. and Barboric, M. (2010) *RNA Biol* 7, 145-50.
5. Pirngruber, J. et al. (2009) *Cell Cycle* 8, 3636-42.
6. Wada, T. et al. (1998) *EMBO J* 17, 7395-403.
7. Yamada, T. et al. (2006) *Mol Cell* 21, 227-37.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **ChIP:** Chromatin IP **C&R:** CUT&RUN

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.