

#68083 Store at -20C

HPC4-Tag Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: All	Sensitivity: Transfected Only	Source: Rabbit
<hr/>			
Product Usage Information	Application		Dilution
	Western Blotting		1:1000
	Immunoprecipitation		1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.		
Specificity / Sensitivity	HPC4-Tag Antibody recognizes recombinant proteins containing the HPC4-tag. This antibody recognizes the HPC4-tag fused to either the amino or carboxy terminus of targeted proteins in transfected cells.		
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues of the HPC4-tag.		
Background	<p>Epitope tags are useful for the labeling and detection of proteins using immunoblotting, immunoprecipitation, and immunostaining techniques. Because of their small size, they are unlikely to affect the tagged protein's biochemical properties.</p> <p>HPC4-tag is a 12 amino acid sequence derived from protein C, which is a vitamin K-dependent plasma zymogen. Protein C is activated by proteolytic cleavage of the thrombin-thrombomodulin complex to form an anticoagulant enzyme (1,2). This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion (3,4).</p>		
Background References	<ol style="list-style-type: none">1. Kurosawa, S. (1989) <i>Nihon Ketsueki Gakkai Zasshi</i> 52, 1343-9.2. Rezaie, A.R. and Esmon, C.T. (1992) <i>J Biol Chem</i> 267, 26104-9.3. Tie, J.K. et al. (2005) <i>J Biol Chem</i> 280, 16410-6.4. Chu, P.H. et al. (2006) <i>Proc Natl Acad Sci U S A</i> 103, 19308-13.		

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the</p>

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.