

#21799 Store at +4°C

Aiolos (D1C1E) Rabbit mAb (PE Conjugate)


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA
For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #Q9UKT9	Entrez-Gene Id: 22806
-------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
Specificity / Sensitivity	Aiolos (D1C1E) Rabbit mAb (PE Conjugate) recognizes endogenous levels of total Aiolos protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro46 of human Aiolos protein.	
Product Description	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Aiolos (D1C1E) Rabbit mAb #15103.	
Background	<p>Aiolos is an Ikaros family transcription factor composed of several zinc fingers that mediate DNA binding and homodimerization or heterodimerization with other Ikaros family members (1). Multiple Aiolos isoforms are generated through alternative splicing of the portion of the transcript encoding the amino-terminal zinc fingers (2). Aiolos is expressed by lymphoid tissues, with highest expression levels seen in mature B and T cells (1). Ikaros family proteins control lymphocyte development by recruiting chromatin remodeling complexes to DNA (3). B cells from mice lacking Aiolos have a reduced threshold for activation, increased proliferation, and elevated levels of IgG and IgE. In addition, Aiolos null mice develop B cell lymphomas (4). In T cells, Aiolos contributes to Th17 cell differentiation by suppressing IL-2 expression (5). Aberrant expression of Aiolos in transformed epithelial cells promotes anchorage independence through downregulation of adhesion-related genes (6). Alterations in the <i>Aiolos</i> gene are observed in near haploid acute lymphoblastic leukemia, and the genetic locus containing Aiolos is linked to increased susceptibility to rheumatoid arthritis and systemic lupus erythematosus (7-9).</p>	
Background References	<ol style="list-style-type: none"> 1. Morgan, B. et al. (1997) <i>EMBO J</i> 16, 2004-13. 2. Liippo, J. et al. (2001) <i>Eur J Immunol</i> 31, 3469-74. 3. Kim, J. et al. (1999) <i>Immunity</i> 10, 345-55. 4. Wang, J.H. et al. (1998) <i>Immunity</i> 9, 543-53. 5. Quintana, F.J. et al. (2012) <i>Nat Immunol</i> 13, 770-7. 6. Li, X. et al. (2014) <i>Cancer Cell</i> 25, 575-89. 7. Holmfeldt, L. et al. (2013) <i>Nat Genet</i> 45, 242-52. 8. Kurreeman, F.A. et al. (2012) <i>Am J Hum Genet</i> 90, 524-32. 9. Lessard, C.J. et al. (2012) <i>Am J Hum Genet</i> 90, 648-60. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	<p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.</p>

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.