e at -20C	SAP102 (A7R8L) Rabbit mAb	T C	Cell Signaling		
Store at		Orders:	877-616-CELL (2355) orders@cellsignal.com		
21		Support:	877-678-TECH (8324)		
474		Web:	info@cellsignal.com cellsignal.com		
#		3 Trask Lane Danvers M	assachusetts 01923 USA		

Applications	Desetivity	Consistivity	N #11A / /
For Research Use On	ly. Not for Use in	Diagnostic Proc	edures.

Applications: WB, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 102	Source/Isotype: Rabbit	UniProt ID: #Q92796	Entrez-Gene Id: 1741	
Product Usage Information	We	plication estern Blotting munoprecipitation			Dilution 1:1000 1:50		
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					erol and less than	
Specificity / Sensitivity		SAP102 (A7R8L) Rabbit mAb recognizes endogenous levels of total SAP102 protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human SAP102 protein.					
Background		Synapse-associated protein 102 (SAP102, DLG3) belongs to the membrane-associated guanylate kinase (MAGUK) protein family and is a homolog of the <i>Drosophila</i> disc large (dlg) tumor suppressor protein. SAP102 consists of three PDZ domains, a Src homology 3 (SH3) domain, and a guanylate kinase (GK) domain (1). The SAP102 protein is more highly expressed in nonproliferating cells than in proliferating cells, indicating a role in the negative regulation of cell growth. SAP102 interacts with the carboxy terminus of the adenomatous polyposis coli (APC) tumor suppressor protein. Furthermore, SAP102 associates with PSD95 in the presence of calcium while the SH3 domain of SAP102 binds calmodulin (2,3). All three PDZ domains of SAP102 participate in binding to the NMDA receptor, interacting specifically with the carboxy-terminal domain of the N-methyl-D-aspartate receptor 2B (NR2B). This SAP102-NR2B interaction may facilitate AMPA receptor withdrawal from the postsynaptic membrane by inhibiting the Erk/MAPK pathway (1,4). Neuronal SAP102 is concentrated at dendritic shafts and spines, axons, and synaptic junctions. At excitatory synapses, SAP102 is involved in NMDA receptor clustering and immobilization and links NMDA receptors to the submembraneous cytomatrix (4). SAP102 and the NMDA receptor function together to mediate plasticity, behavior, and signal transduction (1). A nonsyndromic form of X-linked mental retardation is caused by loss-of-function mutations to the SAP102 gene. The SAP102 protein may be involved in autism since MAGUK proteins in the NMDA receptor complex bind directly to the autism susceptibility gene, neuroligin (1,5).					
Background Refere	2. N 3. N 4. L	Cuthbert, P.C. et al. (Iakino, K. et al. (199 Iasuko, N. et al. (199 au, L.F. et al. (1996) an, J. et al. (2005) M	97) Oncogene 14 99) J Biol Chem J Biol Chem 27:	, 2425-33. 274, 5782-90. I, 21622-8.			
Species Reactivity	Spec	cies reactivity is dete	ermined by testing	g in at least one approve	ed application (e.g., we	estern blot).	
Western Blot Buffe				membrane with diluted th gentle shaking, overr		% w/v nonfat dry	
Applications Key	WB	: Western Blotting IF	P: Immunoprecip	itation			
Cross-Reactivity K	X: X		B: bovine Dg: d	Mk: monkey Vir: virus I og Pg: pig Sc: S. cerevi es expected		0	
Trademarks and Patents				of Cell Signaling Techno aling Technology, Inc.	ology, Inc.		

SAP102 (A7R8L) Rabbit mAb (#47421) Datasheet Without Images Cell Signaling Technology

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose, Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.