

#36057 Store at -20°C

SNAT1/SLC38A1 (D9L2P) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R	Endogenous	50, 70	Rabbit IgG	#Q9H2H9	81539

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

SNAT1/SLC38A1 (D9L2P) Rabbit mAb recognizes endogenous levels of total SNAT1/SLC38A1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly37 of human SNAT1/SLC38A1 protein.

Background

SNAT1/SLC38A1 belongs to the system A transporters that mediate Na⁺-dependent transport of short-chain neutral amino acids such as alanine, serine, and glutamine. SNAT1/SLC38A1 mediates the uptake of glutamine in neurons and plays a crucial role in glutamate-glutamine cycle. Steep concentration gradients across the plasma membrane are achieved by coupling of the electrochemical sodium gradient to amino acid transport. This allows a unidirectional mode of transport for SNAT1/SLC38A1. Upregulation of SNAT1/SLC38A1 by neurotrophic factors is key to dendritic growth and branching of cortical neurons. High expression of SNAT1/SLC38A1 is found in cerebral cortex primarily in neurons and to a lesser extent in astrocytes (1-4). Elevated SNAT1/SLC38A1 expression is prominent in human solid tumors including gliomas, hepatocellular carcinomas and human breast cancer (5-8). Research studies show that an aberrant SNAT1/SLC38A1 expression profile correlates with solid tumor recurrence and poor prognosis in patients with cholangiocarcinoma (9).

Background References

1. Yao, D. et al. (2000) *J Biol Chem* 275, 22790-7.
2. Mackenzie, B. et al. (2003) *J Biol Chem* 278, 23720-30.
3. Chaudhry, F.A. et al. (2002) *J Cell Biol* 157, 349-55.
4. Yu, W.L. et al. (2011) *J Surg Res* 171, 663-8.
5. Melone, M. et al. (2004) *Cereb Cortex* 14, 562-74.
6. Kondoh, N. et al. (2007) *Int J Oncol* 31, 81-7.
7. Sidoryk, M. et al. (2004) *Neuroreport* 15, 575-8.
8. Wang, K. et al. (2013) *BMC Cancer* 13, 343.
9. Burkhalter, J. et al. (2007) *J Biol Chem* 282, 5152-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.