3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: React FC-FP H M		UniProt ID:Entrez-Gene Id:#P434056850
Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide a antibody. Protect from light. Do not freeze.	nd 2 mg/ml BSA. Store at 4°C. Do not aliquot the
Specificity / Sensitivity	Syk (D3Z1E) XP [®] Rabbit mAb (PE Conjugate) recognize	s endogenous levels of total Syk protein.
Source / Purification	Monoclonal antibody is produced by immunizing animals residues around Asn463 of human Syk protein.	with a synthetic peptide corresponding to
Product Description	This Cell Signaling Technology antibody is conjugated to flow cytometry analysis in human cells. This antibody is e reactivity as the unconjugated Syk (D3Z1E) XP [®] Rabbit r	expected to exhibit the same species cross-
Background	Syk is a protein tyrosine kinase that plays an important ro- hematopoietic cells (1-3). Syk interacts with immunorecep- located in the cytoplasmic domains of immune receptors downstream signaling events that mediate diverse cellula differentiation, and phagocytosis (4). There is also eviden investigators have indicated that Syk is a potential tumor Tyr323 is a negative regulatory phosphorylation site withi Phosphorylation at Tyr323 provides a direct binding site f involved in the association of PLCy1 (8). Tyr525 and Tyr5 kinase domain; phosphorylation at Tyr525/526 of human essential for Syk function (9).	ptor tyrosine-based activation motifs (ITAMs) (4). It couples the activated immunoreceptors to ar responses, including proliferation, ace of a role for Syk in nonimmune cells and suppressor in human breast carcinomas (5). In the SH2-kinase linker region in Syk. or the TKB domain of Cbl (6,7). Tyr352 of Syk is 526 are located in the activation loop of the Syk
Background References	 Cheng, A.M. and Chan, A.C. (1997) <i>Curr Opin Immunol</i> Kurosaki, T. (1997) <i>Curr Opin Immunol</i> 9, 309-18. Chu, D.H. et al. (1998) <i>Immunol Rev</i> 165, 167-80. Turner, M. et al. (2000) <i>Immunol Today</i> 21, 148-54. Coopman, P.J. et al. (2000) <i>Nature</i> 406, 742-7. Deckert, M. et al. (1998) <i>J Biol Chem</i> 273, 8867-74. Rao, N. et al. (2001) <i>EMBO J</i> 20, 7085-95. Law, C.L. et al. (1996) <i>Mol Cell Biol</i> 16, 1305-15. Zhang, J. et al. (2000) <i>J Biol Chem</i> 275, 35442-7. 	n/ 9, 528-33.
Species Reactivity	Species reactivity is determined by testing in at least one a	approved application (e.g., western blot).
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)	
Cross-Reactivity Key	 H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S GP: Guinea Pig Rab: rabbit All: all species expected 	-
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling XP is a registered trademark of Cell Signaling Technology, All other trademarks are the property of their respective ov information.	, Inc.

Limited Uses

Syk (D3Z1E) XP® Rabbit mAb (PE Conjugate) (#83324) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.