CD28 (D2Z4E) Rabbit mAb					BISignaling CHNOLOGY® 877-616-CELL (2355) orders@cellsignal.com 877-678-TECH (8324) info@cellsignal.com cellsignal.com
#			3 Trask La	ne Danvers Ma	ssachusetts 01923 USA
For Research Use Only. Not for Use in Diagnostic Procedures.					
Applications: Reactivi WB, IP, IHC-Bond, IHC- H M P, IF-F	ity: Sensitivity: Endogenous	MW (kDa): 14, 40-60	Source/Isotype: Rabbit IgG	UniProt ID: #P10747	Entrez-Gene Id: 940
Product Usage	Application			Dilution	
Information	Western Blotting			1:1000	
	Immunoprecipitation			1:50	
	IHC Leica Bond			1:400 - 1:	1600
	Immunohistochemistry	(Paraffin)		1:50 - 1:2	00
	Immunofluorescence (I	=rozen)		1:50	
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
	For a carrier free (BSA a	and azide free) ve	ersion of this product see	product #87823.	
Specificity / Sensitivity	CD28 (D2Z4E) Rabbit n with multiple CD28 isofc	nAb recognizes e orms that are pro- oteins of unknow	ndogenous levels of total duced through alternative n origin in some mouse ce	CD28 protein and splicing. This anti	body cross-reacts with
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro208 of human CD28 protein.			rresponding to	
Background	CD28 is a transmembrane glycoprotein expressed by T cells as well as some other hematopoietic cells (1,2). T cell activation requires T cell receptor (TCR) recognition of antigen presented in the context of MHC molecules. CD28 acts as a T cell costimulatory receptor, and interaction of CD28 with its ligands CD80 or CD86 provides the second signal required for naïve T cell activation (3-5). Activation of naïve T cells in the absence of CD28 stimulation can result in a state of T cell anergy, or unresponsiveness (3). CD28 signals through cytoplasmic phospho-tyrosine motifs that bind several SH2 or SH3 domain-containing proteins involved in T cell activation (2). Recently, CD28 was demonstrated to be a preferred target of PD-1-mediated dephosphorylation. Consistently, CD28 expression was required for T cell proliferation following PD-1 blockade and CD28 stimulation was required for effective anti-PD-1 cancer immunotherapy in mice (6,7). Several CD28 isoforms are produced by alternative splicing (8).				
Background References	 Aruffo, A. and Seed, B. (1987) <i>Proc Natl Acad Sci U S A</i> 84, 8573-7. Esensten, J.H. et al. (2016) <i>Immunity</i> 44, 973-88. Harding, F.A. et al. (1992) <i>Nature</i> 356, 607-9. Azuma, M. et al. (1993) <i>Nature</i> 366, 76-9. Linsley, P.S. et al. (1990) <i>Proc Natl Acad Sci U S A</i> 87, 5031-5. Hui, E. et al. (2017) <i>Science</i> 355, 1428-1433. Kamphorst, A.O. et al. (2017) <i>Science</i> 355, 1423-1427. Magistrelli, G. et al. (1999) <i>Biochem Biophys Res Commun</i> 259, 34-7. 				
Species Reactivity	Species reactivity is dete	rmined by testing	g in at least one approved	application (e.g.,	western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key	÷		tation IHC-Bond: IHC Lei I F-F: Immunofluorescence		

3/23/24, 11:04 AM Cross-Reactivity Key	CD28 (D2Z4E) Rabbit mAb (#38774) Datasheet Without Images Cell Signaling Technology H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for reseale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.