#64616 Store at -20C

Phospho-FoxO3a (Ser425) (D5E9T) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 82-97	Source/Isotype: Rabbit IgG	UniProt ID: #O43524	Entrez-Gene Id 2309	
Product Usage Information	Ap	plication		Dilution			
	We	stern Blotting		1:1000			
	lmı	munoprecipitation		1:50			
Storage	•	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20 °C. Do not aliquot the antibody.					
Specificity / Sensiti	whe	Phospho-FoxO3a (Ser425) (D5E9T) Rabbit mAb recognizes endogenous levels of FoxO3a protein only when phosphorylated at Ser425. This antibody does not cross-react with FoxO1 nor FoxO4 proteins. This antibody may detect a non-specific band at 65 kDa and at 240 kDa.					
Source / Purificatio		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser425 of human FoxO3a protein.					

The Forkhead family of transcription factors is involved in tumorigenesis of rhabdomyosarcoma and acute leukemias (1-3). Within the family, three members (FoxO1, FoxO4, and FoxO3a) have sequence similarity to the nematode orthologue DAF-16, which mediates signaling via a pathway involving IGFR1, PI3K, and Akt (4-6). Active forkhead members act as tumor suppressors by promoting cell cycle arrest and apoptosis. Increased expression of any FoxO member results in the activation of the cell cycle inhibitor p27 Kip1. Forkhead transcription factors also play a part in TGF-β-mediated upregulation of p21 Cip1, a process negatively regulated through PI3K (7). Increased proliferation results when forkhead transcription factors are inactivated through phosphorylation by Akt at Thr24, Ser256, and Ser319, which results in nuclear export and inhibition of transcription factor activity (8). Forkhead transcription factors can also be inhibited by the deacetylase sirtuin (SirT1) (9).

ERK phosphorylates FoxO3a at Ser425, degrading FoxO3a via the ubiquitin-proteasome pathway. Research has shown that this phosphorylation event inhibits FoxO3a's function to induce cell death and thus promotes cell proliferation and tumorigenesis (10, 11).

Background References

Background

- 1. Anderson, M.J. et al. (1998) Genomics 47, 187-99.
- 2. Galili, N. et al. (1993) Nat Genet 5, 230-5.
- 3. Borkhardt, A. et al. (1997) Oncogene 14, 195-202.
- 4. Nakae, J. et al. (1999) *J Biol Chem* 274, 15982-5.
- 5. Rena, G. et al. (1999) J Biol Chem 274, 17179-83.
- 6. Guo, S. et al. (1999) J Biol Chem 274, 17184-92.
- 7. Seoane, J. et al. (2004) Cell 117, 211-23.
- 8. Arden, K.C. (2004) Mol Cell 14, 416-8.
- 9. Yang, Y. et al. (2005) EMBO J 24, 1021-32.
- 10. Brunet, A. et al. (2004) Science 303, 2011-5.
- 11. Yang, J.Y. et al. (2008) Nat Cell Biol 10, 138-48.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

1/1/24. 12:25 PM

Phospho-FoxO3a (Ser425) (D5E9T) Rabbit mAb (#64616) Datasheet Without Images Cell Signaling Techno...

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dq: dog Pq: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.