3/23/24, 1:11 PM Revision 6

CXCR5 (D6L3C) Rabbit mAb						
Store				Orders:	877-616-CELL (2355) orders@cellsignal.com	
172				Support:	877-678-TECH (8324)	
#7217				Web:	info@cellsignal.com cellsignal.com	
#				3 Trask Lane Danvers	Massachusetts 01923 USA	
For Research Use Only. Not for Use in Diagnostic Procedures.						
Applications: IHC-Bond, IHC-P	Reactivity: H	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P32302	Entrez-Gene Id: 643	

0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier-free (BSA and azide free) version of this product see product #99825 Specificity / Sensitivity CXCR5 (D6L3C) Rabbit mAb recognizes endogenous levels of total CXCR5 protein. Source / Purification Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the extracellular domains of human CXCR5 protein. Background CXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary breat tumors, in correlation with their propensity to grow and metastasize (4). Background References 1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93.	43				
InformationHC Leica Bond1:50 - 1:200Immunohistochemistry (Paraffin)1:200StorageSupplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier-free (BSA and azide free) version of this product see product #99825Specificity / SensitivityCXCR5 (D6L3C) Rabbit mAb recognizes endogenous levels of total CXCR5 protein.Source / PurificationMonoclonal antibody is produced by immunizing animals with recombinant protein specific to the extracellular domains of human CXCR5 protein.BackgroundCXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary breat tumors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfield, D. et al. (2012) <i>Clin Breast Cancer</i> 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).					
StorageSupplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier-free (BSA and azide free) version of this product see product #99825Specificity / Sensitivity Source / PurificationCXCR5 (D6L3C) Rabbit mAb recognizes endogenous levels of total CXCR5 protein. Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the extracellular domains of human CXCR5 protein.BackgroundCXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL3 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary bre turnors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2012) Clin Breast Cancer 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).					
0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier-free (BSA and azide free) version of this product see product #99825 Specificity / Sensitivity CXCR5 (D6L3C) Rabbit mAb recognizes endogenous levels of total CXCR5 protein. Source / Purification Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the extracellular domains of human CXCR5 protein. Background CXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary breat tumors, in correlation with their propensity to grow and metastasize (4). Background References 1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93. Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Specificity / Sensitivity Source / PurificationCXCR5 (D6L3C) Rabbit mAb recognizes endogenous levels of total CXCR5 protein.BackgroundCXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary bre tumors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) Immunity 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Source / PurificationMonoclonal antibody is produced by immunizing animals with recombinant protein specific to the extracellular domains of human CXCR5 protein.BackgroundCXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary bre tumors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).					
BackgroundCXCR5 is a G protein-coupled receptor belonging to the chemokine receptor subfamily (1). Upon bi of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary breat tumors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).					
of its ligand, the chemokine CXCL13, CXCR5 initiates multiple intracellular signaling pathways that regulate cell proliferation, survival, and migration (2). CXCR5 is expressed in both mature B cells ar follicular helper T cells, and responds to CXCL13 gradient to control lymphocyte migration towards secondary lymphoid tissues (3). CXCR5 has also been shown to be highly expressed in primary bre tumors, in correlation with their propensity to grow and metastasize (4).Background References1. Zlotnik, A. and Yoshie, O. (2000) <i>Immunity</i> 12, 121-7. 2. Luster, A.D. (1998) <i>N Engl J Med</i> 338, 436-45. 3. Breitfeld, D. et al. (2000) <i>J Exp Med</i> 192, 1545-52. 4. Razis, E. et al. (2012) <i>Clin Breast Cancer</i> 12, 183-93.Species ReactivitySpecies reactivity is determined by testing in at least one approved application (e.g., western blot).					
2. Luster, A.D. (1998) N Engl J Med 338, 436-45. 3. Breitfeld, D. et al. (2000) J Exp Med 192, 1545-52. 4. Razis, E. et al. (2012) Clin Breast Cancer 12, 183-93. Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).	d				
Applications Key IHC-Bond: IHC Leica Bond IHC-P: Immunohistochemistry (Paraffin)					
Cross-Reactivity KeyH: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melano X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected	Jaster				
Trademarks and PatentsCell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SignalStain is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for r information.	iore				
following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's ter	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purp Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any m that conflicts with its labeling statement. Products sold or licensed by CST are provided for Custome end-user and solely for research and development uses. Any use of Product for diagnostic, prophyla therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other com purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or	anner as the stic or				

CXCR5 (D6L3C) Rabbit mAb (#72172) Datasheet Without Images Cell Signaling Technology

otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.