| Revision 1                        |   |  |  |   |  |   |
|-----------------------------------|---|--|--|---|--|---|
| TRAP1/HSP75 (D3D7N) Rabbit<br>mAb |   |  |  |   |  |   |
| Store                             |   |  |  |   | Orders:  | 877-616-CELL (2355)<br>orders@cellsignal.com  |
| 345                               |   |  |  |   | Support:   | 877-678-TECH (8324)   |
| 929                               |   |  |  |   | Web:   | info@cellsignal.com<br>cellsignal.com   |
| #                                 |   |  |  | 3 Trask L   | .ane   Danvers   Ma  | assachusetts   01923   USA  |
| For Research Use On               | ly. Not for Use in  | Diagnostic Proc  | edures.  |   |  |   |
| Applications:<br>WB, IP, IF-IC    | Reactivity:<br>H  | Sensitivity:<br>Endogenous   | <b>MW (kDa):</b><br>75   | Source/Isotype:<br>Rabbit IgG   | UniProt ID:<br>#Q12931   | Entrez-Gene Id:<br>10131  |
| Product Usage<br>Information      | Арј   | plication  |  |   |  | Dilution  |
|                                   | We  | stern Blotting   |  |   |  | 1:1000  |
|                                   | Imn   | nunoprecipitation  |  |   |  | 1:50  |
|                                   | Imn   | Immunofluorescence (Immunocytochemistry)   |  |   |  | 1:100   |
|                                   |   | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.  |  |   |  |   |
| Specificity / Sens                | sitivity TRA  | TRAP1/HSP75 (D3D7N) Rabbit mAb recognizes endogenous levels of total TRAP1/HSP75 protein.  |  |   |  |   |
| Source / Purifica                 |   | Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro70 of human TRAP1/HSP75 protein.   |  |   |  |   |
| Background                        | ATP<br>of th<br>injur<br>TRA<br>prote<br>cytor<br>cell o<br>mito<br>early | ase that was origin<br>e HSP90 family, TF<br>y, and UV irradiatio<br>P1 within mitochor<br>ective role in ischer<br>chrome C release a<br>death by inhibiting<br>chondrial serine/thu<br>y-onset Parkinson's | ally identified as a<br>RAP1 is not heat-<br>in. An amino-term<br>idria (1). Overexp<br>mia injury (2). Res<br>and apoptosis, wi<br>the generation of<br>reonine kinase PI<br>disease (PD). PI | P1), also known as HSF<br>a protein that interacts w<br>inducible but is upregula<br>inal mitochondrial locali<br>pression of TRAP1 decre<br>search studies demonstr<br>th additional evidence in<br>reactive oxygen species<br>NK1, whose correspond<br>INK1 protects cells from<br>predice ondria BD liab | with the TNF receptor<br>ated by glucose dep<br>zation sequence re<br>eases oxidative stree<br>rate that silencing of<br>indicating that TRAP<br>s (3). TRAP1 is a su<br>ling gene is mutate<br>oxidative stress-ind | or. Although a member<br>privation, oxidative<br>sults in localization of<br>ess, suggesting a<br>of TRAP1 enhances<br>1 can protect cells from<br>ubstrate of the<br>d in some forms of<br>duced cell death by |

suppressing release of cytochrome C from mitochondria. PD-linked *PINK1* mutations impair the ability of PINK1 to phosphorylate TRAP1 and leads to impaired cell survival (4). Finally, TRAP1 alleviates α-synuclein induced toxicity and rescues the PINK1 loss-of-function phenotype (5).
 1. Felts, S.J. et al. (2000) *J Biol Chem* 275, 3305-12.

 Background References
 1. Felts, S.J. et al. (2000) J Biol Chem 275, 3305-12.

 2. Hua, G. et al. (2007) J Biol Chem 282, 20553-60.

 3. Voloboueva, L.A. et al. (2008) J Cereb Blood Flow Metab 28, 1009-16.

 4. Pridgeon, J.W. et al. (2007) PLoS Biol 5, e172.

 5. Butler, E.K. et al. (2012) PLoS Genet 8, e1002488.

| Species Reactivity        | Species reactivity is determined by testing in at least one approved application (e.g., western blot).   |
|---------------------------|--|
| Western Blot Buffer       | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.   |
| Applications Key          | WB: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry)   |
| Cross-Reactivity Key      | <ul> <li>H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster</li> <li>X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse</li> <li>GP: Guinea Pig Rab: rabbit All: all species expected</li> </ul> |
| Trademarks and<br>Patents | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.<br>All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more<br>information.  |
|                           |  |

TRAP1/HSP75 (D3D7N) Rabbit mAb (#92345) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.