

#79299 Store at -20C

Angiotensinogen Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M	Endogenous	55, 70	Rabbit	#P01019	183

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Angiotensinogen Antibody recognizes endogenous levels of total angiotensinogen protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of mouse angiotensinogen protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Angiotensinogen (AGT) is the primary precursor of angiotensins, peptide hormones that play a central role in the renin-angiotensin system (RAS) (1-3). AGT is a secreted protein synthesized primarily by the liver and secreted into circulation. Upon binding to renin, the amino terminal fragment of AGT is cleaved and released as a decapeptide hormone termed angiotensin 1 (Ang I). Ang I is subsequently processed by angiotensin converting enzyme (ACE) to generate angiotensin II (Ang II), which acts on AT1 and AT2 receptors in the central nervous system to increase production of anti-diuretic hormone (ADH), while promoting vasoconstriction in the peripheral circulation (4). Aberrant upregulation of Ang II has been associated with numerous clinical conditions, including hypertension, atherosclerosis, myocardial hypertrophy, and obesity (5-7). Alternative cleavage products of Ang I (e.g., Ang 1-7) can also be generated by ACE2 cleavage, some of which display biological functions that are distinct from Ang II (8). Treatments that target the RAS (e.g., ACE inhibitors) are consequently of significant importance in the treatment of hypertensive and hypertensive-related disorders(5-8).

Background References

1. Lu, H. et al. (2016) *Hypertens Res* 39, 492-500.
2. Sparks, M.A. et al. (2014) *Compr Physiol* 4, 1201-28.
3. Kumar, R. et al. (2012) *Clin Sci (Lond)* 123, 273-84.
4. de Kloet, A.D. et al. (2015) *Am J Physiol Regul Integr Comp Physiol* 309, R444-58.
5. Vajapey, R. et al. (2014) *Front Physiol* 5, 439.
6. Wu, C. et al. (2011) *N Am J Med Sci (Boston)* 4, 183-190.
7. Putnam, K. et al. (2012) *Am J Physiol Heart Circ Physiol* 302, H1219-30.
8. Padda, R.S. et al. (2015) *J Diabetes Metab* 6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.