

#59995 Store at -20°C

c-Myb (D1B9E) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IHC-P, FC-FP	H	Endogenous	80	Rabbit IgG	#P10242	4602

Product Usage Information

Application

Western Blotting
Immunohistochemistry (Paraffin)
Flow Cytometry (Fixed/Permeabilized)

Dilution

1:1000
1:200
1:400

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

c-Myb (D1B9E) Rabbit mAb recognizes endogenous levels of total c-Myb protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human c-Myb protein.

Background

c-Myb is a transcriptional activator that specifically recognizes the sequence 5'-YAAC[GT]G-3'. It is expressed in hematopoietic progenitor cells where it plays an important role in the control of proliferation and differentiation (1-3). c-Myb is required for transcription of genes involved in self-renewal of intestinal stem cells. Importantly, c-Myb regulates expression of Lgr5, a protein expressed in putative intestinal stem cells that give rise to all cell lineages of small intestinal crypts (4). c-Myb is reported to be expressed in colon crypt cells and in human colorectal cancer lines (5,6). Research has shown that c-Myb gene translocations and copy number alterations are found in several leukemias, breast cancer, and other solid tumors (7,8).

Background References

1. Lin, H.H. et al. (1996) *Curr Top Microbiol Immunol* 211, 79-87.
2. Mucenski, M.L. et al. (1991) *Cell* 65, 677-89.
3. Badiani, P. et al. (1994) *Genes Dev* 8, 770-82.
4. Cheasley, D. et al. (2011) *Stem Cells* 29, 2042-50.
5. Thompson, M.A. et al. (1998) *Cancer Res* 58, 5168-75.
6. Wilkins, H.R. et al. (2010) *Tumour Biol* 31, 16-22.
7. Ramsay, R.G. and Gonda, T.J. (2008) *Nat Rev Cancer* 8, 523-34.
8. Stenman, G. et al. (2010) *Cell Cycle* 9, 2986-95.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IHC-P:** Immunohistochemistry (Paraffin) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
Alexa Fluor is a registered trademark of Life Technologies Corporation.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.