

#99902 Store at +4°C

Phospho-Stat6 (Tyr641) (D8S9Y) Rabbit mAb (PE Conjugate)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P42226	Entrez-Gene Id: 6778
-------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
Specificity / Sensitivity	Phospho-Stat6 (Tyr641) (D8S9Y) Rabbit mAb (PE Conjugate) recognizes endogenous levels of Stat6 protein only when phosphorylated at Tyr641. Weak cross-reactivity with other tyrosine-phosphorylated proteins has been observed in some cell lines.	
Species predicted to react based on 100% sequence homology:	Rat	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic phospho-peptide corresponding to residues surrounding Tyr641 of human Stat6 protein.	
Product Description	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-Stat6 (Tyr641) (D8S9Y) Rabbit mAb #56554.	
Background	Upon activation by Janus kinases, Stat6 translocates to the nucleus where it regulates cytokine-induced gene expression. Stat6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13 (1-4). In addition, Stat6 is activated by IFN- α in B cells, where it forms transcriptionally active complexes with Stat2 and p48 (5,6). Protein phosphatase 2A is also involved in regulation of IL-4-mediated Stat6 signaling (7).	
Background References	<ol style="list-style-type: none"> Nelms, K. et al. (1999) <i>Ann. Rev. Immunol.</i> 17, 701-738. Malabarba, M.G. et al. (1996) <i>Biochem. J.</i> 319, 865-872. Hou, J. et al. (1994) <i>Science</i> 265, 1701-1706. Quelle, F.W. et al. (1995) <i>Mol. Cell. Biol.</i> 15, 3336-3343. Takeda, K. et al. (1996) <i>Nature</i> 380, 627-630. Gupta, S. et al. (1999) <i>J. Immunol.</i> 163, 3834-3841. Woetmann, A. et al. (2003) <i>J. Biol. Chem.</i> 278, 2787-2791. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.