

#57981 Store at -20°C

**FBXO7 (D8L4E) Rabbit mAb****Cell Signaling**  
TECHNOLOGY®**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M	Endogenous	65-72	Rabbit IgG	#Q9Y3I1	25793

<b>Product Usage Information</b>	<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	FBXO7 (D8L4E) Rabbit mAb recognizes endogenous levels of total FBXO7 protein. FBXO7 (D8L4E) Rabbit mAb detects two bands in human cerebellum, consistent with multiple FBXO7 isoforms generated by alternative splicing.	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the N-terminus of human FBXO7 protein.	
<b>Background</b>	F-box only protein 7 (FBXO7) is the substrate recognition component of the SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex. This complex mediates ubiquitination and degradation of targeted proteins. Several "PARK" chromosomal loci have been associated with monogenic forms of PD, including <i>PARK15</i> (1). The associated allele for <i>PARK15</i> is <i>FBXO7</i> , which encodes the FBXO7 protein. Mutations in <i>FBXO7</i> cause early-onset autosomal recessive PD, and alterations in FBXO7 normal function, e.g. the protein degradation process, may contribute to PD etiology (2, 3). Additionally, expression of specific FBXO7 isoforms generated by alternative splicing is linked with PD (4).	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Walden, H. and Muqit, M.M. (2017) <i>Biochem J</i> 474, 1439-1451.</li> <li>2. Burchell, V.S. et al. (2013) <i>Nat Neurosci</i> 16, 1257-65.</li> <li>3. Vingill, S. et al. (2016) <i>EMBO J</i> 35, 2008-25.</li> <li>4. Zhao, T. et al. (2011) <i>PLoS One</i> 6, e16983.</li> <li>5. Zhao, T. et al. (2011) <i>PLoS One</i> 6, e16983.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>WB:</b> Western Blotting
<b>Cross-Reactivity Key</b>	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or</p>

therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.