

#98110 Store at -20°C

## Necroptosis Antibody Sampler Kit



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

1 Kit (6 x 20 microliters)

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
RIP (D94C12) XP® Rabbit mAb	3493	20 µl	78 kDa	Rabbit IgG
Phospho-RIP (Ser166) (D1L3S) Rabbit mAb	65746	20 µl	78-82 kDa	Rabbit IgG
MLKL (D2I6N) Rabbit mAb	14993	20 µl	54 kDa	Rabbit IgG
Phospho-MLKL (Ser358) (D6H3V) Rabbit mAb	91689	20 µl	54 kDa	Rabbit IgG
RIP3 (E1Z1D) Rabbit mAb	13526	20 µl	46-62 kDa	Rabbit IgG
Phospho-RIP3 (Ser227) (D6W2T) Rabbit mAb	93654	20 µl	46-62 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit [cellsignal.com](http://cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

### Description

The Necroptosis Antibody Sampler Kit provides an economical means of detecting total and phosphorylated proteins associated with necroptosis. The kit includes enough antibody to perform two western blots with each primary antibody.

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Background

Necroptosis, a regulated pathway for necrotic cell death, is triggered by a number of inflammatory signals, including cytokines in the tumor necrosis factor (TNF) family, pathogen sensors such as toll-like receptors (TLRs), ischemic injury, and neurodegenerative diseases (1-3). The process is negatively regulated by caspases and is initiated through a complex containing the RIP and RIP3 kinases, typically referred to as the necrosome. Necroptosis is inhibited by a small molecule inhibitor of RIP, necrostatin-1 (Nec-1) (4). RIP is phosphorylated at several sites within the kinase domain that are sensitive to Nec-1, including Ser14, Ser15, Ser161, and Ser166 (5). During necroptosis, RIP3 is phosphorylated at Ser227, leading to recruitment and phosphorylation of MLKL at Thr357 and Ser358 (6). Phosphorylation of MLKL results in its oligomerization and translocation to the plasma membrane, where it affects membrane integrity (7-10).

### Background References

1. Christofferson, D.E. and Yuan, J. (2010) *Curr Opin Cell Biol* 22, 263-8.
2. Kaczmarek, A. et al. (2013) *Immunity* 38, 209-23.
3. Zhou, W. and Yuan, J. (2014) *Semin Cell Dev Biol* 35, 14-23.
4. Degterev, A. et al. (2008) *Nat Chem Biol* 4, 313-21.
5. Ofengeim, D. and Yuan, J. (2013) *Nat Rev Mol Cell Biol* 14, 727-36.
6. Sun, L. et al. (2012) *Cell* 148, 213-27.
7. Cai, Z. et al. (2014) *Nat Cell Biol* 16, 55-65.
8. Chen, X. et al. (2014) *Cell Res* 24, 105-21.
9. Wang, H. et al. (2014) *Mol Cell* 54, 133-46.
10. Dondelinger, Y. et al. (2014) *Cell Rep* 7, 971-81.

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
XP is a registered trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner

that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.