

#34696 Store at -20°C

TrK (pan) (A7H6R) Rabbit mAb (Biotinylated)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	120-140	Rabbit IgG	#P04629, #Q16288, #Q16620	4914, 4916, 4915

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
Storage	Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Trk (pan) (A7H6R) Rabbit mAb (Biotinylated) detects endogenous levels of total Trk protein. This antibody detects TrkA, TrkB and TrkC. However, the antibody may preferentially detect TrkA over TrkB and TrkC.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide surrounding Tyr791 of human TrkA.	
Product Description	This Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated antibody is expected to exhibit the same species cross-reactivity as the unconjugated Trk (pan) (A7H6R) Rabbit mAb #92991.	

MW (kDa)	120-140
----------	---------

Background	The family of Trk receptor tyrosine kinases consists of TrkA, TrkB, and TrkC. While the sequence of these family members is highly conserved, they are activated by different neurotrophins: TrkA by NGF, TrkB by BDNF or NT4, and TrkC by NT3 (1). Neurotrophin signaling through these receptors regulates a number of physiological processes, such as cell survival, proliferation, neural development, and axon and dendrite growth and patterning (1). In the adult nervous system, the Trk receptors regulate synaptic strength and plasticity. TrkA regulates proliferation and is important for development and maturation of the nervous system (2). Phosphorylation at Tyr490 is required for Shc association and activation of the Ras-MAP kinase cascade (3,4). Residues Tyr674/675 lie within the catalytic domain, and phosphorylation at these sites reflects TrkA kinase activity (3-6). Point mutations, deletions, and chromosomal rearrangements (chimeras) cause ligand-independent receptor dimerization and activation of TrkA (7-10). TrkA is activated in many malignancies including breast, ovarian, prostate, and thyroid carcinomas (8-13). Research studies suggest that expression of TrkA in neuroblastomas may be a good prognostic marker as TrkA signals growth arrest and differentiation of cells originating from the neural crest (10).
-------------------	--

Background References	<ol style="list-style-type: none"> Huang, E.J. and Reichardt, L.F. (2003) <i>Annu Rev Biochem</i> 72, 609-42. Segal, R.A. and Greenberg, M.E. (1996) <i>Annu Rev Neurosci</i> 19, 463-89. Stephens, R.M. et al. (1994) <i>Neuron</i> 12, 691-705. Marsh, H.N. et al. (2003) <i>J Cell Biol</i> 163, 999-1010. Obermeier, A. et al. (1993) <i>EMBO J</i> 12, 933-41. Obermeier, A. et al. (1994) <i>EMBO J</i> 13, 1585-90. Arevalo, J.C. et al. (2001) <i>Oncogene</i> 20, 1229-34. Reuther, G.W. et al. (2000) <i>Mol Cell Biol</i> 20, 8655-66. Greco, A. et al. (1997) <i>Genes Chromosomes Cancer</i> 19, 112-23. Pierotti, M.A. and Greco, A. (2006) <i>Cancer Lett</i> 232, 90-8. Lagadec, C. et al. (2009) <i>Oncogene</i> 28, 1960-70. Greco, A. et al. (2010) <i>Mol Cell Endocrinol</i> 321, 44-9. Ødegaard, E. et al. (2007) <i>Hum Pathol</i> 38, 140-6.
------------------------------	---

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
---------------------------	--

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.