

#17229 Store at -20°C

Pro-Survival Bcl-2 Family Antibody Sampler Kit II

1 Kit (7 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Bcl-2 (D55G8) Rabbit mAb	4223	20 µl	26 kDa	Rabbit IgG
Phospho-Bcl-2 (Ser70) (5H2) Rabbit mAb	2827	20 µl	28 kDa	Rabbit IgG
Mcl-1 (D2W9E) Rabbit mAb	94296	20 µl	40 (human), 35 (rodent) kDa	Rabbit IgG
Phospho-Mcl-1 (Thr163) (D5M9D) Rabbit mAb	14765	20 µl	40 kDa	Rabbit IgG
Bcl-xL (54H6) Rabbit mAb	2764	20 µl	30 kDa	Rabbit IgG
A1/Bfl-1 (D1A1C) Rabbit mAb	14093	20 µl	18 kDa	Rabbit IgG
Bcl-w (31H4) Rabbit mAb	2724	20 µl	18 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Pro-Survival Bcl-2 Family Antibody Sampler Kit II provides an economical means to examine several members of the Bcl-2 family. The kit contains enough primary antibody to perform two western blot experiments.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

The Bcl-2 family consists of a number of evolutionarily conserved proteins containing Bcl-2 homology domains (BH) that regulate apoptosis through control of mitochondrial membrane permeability and release of cytochrome c (1-3). Four BH domains have been identified (BH1-4) that mediate protein interactions. The family can be separated into three groups based upon function and sequence homology: pro-survival members include Bcl-2, Bcl-xL, Mcl-1, A1 and Bcl-w; pro-apoptotic proteins include Bax, Bak and Bok; and "BH3 only" proteins Bad, Bik, Bid, Puma, Bim, Bmf, Noxa and Hrk. Interactions between death-promoting and death-suppressing Bcl-2 family members has led to a rheostat model in which the ratio of pro-apoptotic and anti-apoptotic proteins controls cell fate (4). Thus, pro-survival members exert their behavior by binding to and antagonizing death-promoting members. In general, the "BH3-only members" can bind to and antagonize the pro-survival proteins leading to increased apoptosis (5). While some redundancy of this system likely exists, tissue specificity, transcriptional and post-translational regulation of many of these family members can account for distinct physiological roles.

Several phosphorylation sites have been identified within Bcl-2 including Thr56, Ser70, Thr74 and Ser87 (6). These phosphorylation sites may be targets of the ASK1/MKK7/JNK1 pathway, and phosphorylation of Bcl-2 may be a marker for mitotic events (7,8). Mutation of Bcl-2 at Thr56 or Ser87 inhibits its anti-apoptotic activity during glucocorticoid-induced apoptosis of T lymphocytes (9). Interleukin 3 and JNK-induced Bcl-2 phosphorylation at Ser70 may be required for its enhanced antiapoptotic functions (10). Mcl-1 is phosphorylated in response to treatment with phorbol ester, microtubule-damaging agents, oxidative stress, and cytokine withdrawal (11-14). Phosphorylation at Thr163, the conserved MAP kinase/ERK site located within the PEST region, slows Mcl-1 protein turnover (13) but may prime the GSK-3 mediated phosphorylation at Ser159 that leads to Mcl-1 destabilization (14).

Background References

1. Cory, S. et al. (2003) *Oncogene* 22, 8590-607.
2. Antonsson, B. and Martinou, J.C. (2000) *Exp Cell Res* 256, 50-7.
3. Sharpe, J.C. et al. (2004) *Biochim Biophys Acta* 1644, 107-13.
4. Korsmeyer, S.J. et al. (1993) *Semin Cancer Biol* 4, 327-32.
5. Bouillet, P. and Strasser, A. (2002) *J Cell Sci* 115, 1567-74.
6. Maundrell, K. et al. (1997) *J Biol Chem* 272, 25238-42.
7. Yamamoto, K. et al. (1999) *Mol Cell Biol* 19, 8469-78.
8. Ling, Y.H. et al. (1998) *J Biol Chem* 273, 18984-91.
9. Huang, S.T. and Cidlowski, J.A. (2002) *FASEB J* 16, 825-32.
10. Deng, X. et al. (2001) *J Biol Chem* 276, 23681-8.
11. Domina, A.M. et al. (2000) *J Biol Chem* 275, 21688-94.

12. Inoshita, S. et al. (2002) *J Biol Chem* 277, 43730-4.
13. Domina, A.M. et al. (2004) *Oncogene* 23, 5301-15.
14. Maurer, U. et al. (2006) *Mol Cell* 21, 749-60.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
SignalFire is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.