#98322 Store at -20°C

Pro-Apoptosis Bcl-2 Family Antibody Sampler Kit II



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

.

Web:

info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

1 Kit (9 x 20 microliters)

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source	
Bax (D2E11) Rabbit mAb	5023	20 μΙ	20 kDa	Rabbit IgG	
Bak (D4E4) Rabbit mAb	12105	20 μΙ	25 kDa	Rabbit IgG	
Bok (D7V2N) Rabbit mAb	86875	20 μΙ	22 kDa	Rabbit IgG	
Bim (C34C5) Rabbit mAb	2933	20 μΙ	12, 15, 23 kDa	Rabbit IgG	
Bad (D24A9) Rabbit mAb	9239	20 μΙ	23 kDa	Rabbit IgG	
BID Antibody	2002	20 μΙ	15, 22 kDa	Rabbit	
Puma (D30C10) Rabbit mAb	12450	20 μΙ	23 kDa	Rabbit IgG	
Noxa (D8L7U) Rabbit mAb	14766	20 μΙ	10 kDa	Rabbit IgG	
Bik Antibody	4592	20 μΙ	20 kDa	Rabbit	
Anti-rabbit IgG, HRP-linked Antibody	7074	100 μΙ		Goat	

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Pro-Apoptosis Bcl-2 Family Antibody Sampler Kit II provides an economical means to examine several members of the Bcl-2 family. The kit contains enough primary antibody to perform two western blot experiments.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibodies.

Background

The Bcl-2 family consists of a number of evolutionarily conserved proteins containing Bcl-2 homology domains (BH) that regulate apoptosis through control of mitochondrial membrane permeability and release of cytochrome c (1-3). Four BH domains have been identified (BH1-4) that mediate protein interactions. The family can be separated into three groups based upon function and sequence homology: pro-survival members include Bcl-2, Bcl-xL, Mcl-1, A1 and Bcl-w; pro-apoptotic proteins include Bax, Bak and Bok; and "BH3 only" proteins Bad, Bik, Bid, Puma, Bim, Bmf, Noxa and Hrk. Interactions between death-promoting and death-suppressing Bcl-2 family members has led to a rheostat model in which the ratio of pro-apoptotic and anti-apoptotic proteins controls cell fate (4). Thus, pro-survival members exert their behavior by binding to and antagonizing death-promoting members. In general, the "BH3-only members" can bind to and antagonize the pro-survival proteins leading to increased apoptosis (5). While some redundancy of this system likely exists, tissue specificity, transcriptional and post-translational regulation of many of these family members can account for distinct physiological roles.

Background References

- 1. Cory, S. et al. (2003) Oncogene 22, 8590-607.
- 2. Antonsson, B. and Martinou, J.C. (2000) Exp Cell Res 256, 50-7.
- 3. Sharpe, J.C. et al. (2004) Biochim Biophys Acta 1644, 107-13.
- 4. Korsmeyer, S.J. et al. (1993) Semin Cancer Biol 4, 327-32.
- 5. Bouillet, P. and Strasser, A. (2002) J Cell Sci 115, 1567-74.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.