#68978 store at -20C

MAGE-A10 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 62	Source: Rabbit	UniProt ID: #P43363	Entrez-Gene Id: 4109	
Product Usage Information	Ар	Application			Dilution		
	We	Western Blotting			1:1000		
	Imr	Immunoprecipitation			1:100		
Storage	•	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensitiv		MAGE-A10 Antibody recognizes endogenous levels of total MAGE-A10 protein. This antibody does not cross-react with other MAGE-A family members.					
Source / Purification	resid	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro323 of human MAGE-A10 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background	Can	Cancer/testis antigens (CTAs) are a family of more than 100 proteins whose normal expression is largely					

Cancer/testis antigens (CTAs) are a family of more than 100 proteins whose normal expression is largely restricted to immune privileged germ cells of the testis, ovary, and trophoblast cells of the placenta. Although most normal somatic tissues are void of CTA expression, due to epigenetic silencing of gene expression, their expression is upregulated in a wide variety of human solid and liquid tumors (1,2). As such, CTAs have garnered much attention as attractive targets for a variety of immunotherapy-based approaches to selectively attack tumors (3).

Melanoma-associated antigen-A10 (MAGE-A10) is a cancer testis antigen and belongs to the type I MAGE family of proteins. Unlike other members of the MAGE-A subfamily, MAGE-A10 is a nuclear protein (4). Research studies have shown that MAGE-A10 expression is normally restricted to the human testis and placenta but is aberrantly upregulated in a number of human solid tumors, such as lung cancer, melanoma, bladder cancer, and esophageal carcinoma (4-6). Due to its upregulated expression in human tumors and high degree of immunogenicity, MAGE-A10 has received significant attention as a novel immunotherapy target through the use of vaccines and adoptive cell therapy (7-9).

Background References

- 1. Caballero, O.L. and Chen, Y.T. (2009) Cancer Sci 100, 2014-21.
- 2. De Smet, C. et al. (1999) Mol Cell Biol 19, 7327-35.
- 3. Gjerstorff, M.F. et al. (2015) *Oncotarget* 6, 15772-87.
- 4. Rimoldi, D. et al. (1999) Int J Cancer 82, 901-7.
- 5. Schultz-Thater, E. et al. (2011) Int J Cancer 129, 1137-48.
- 6. Lin, J. et al. (2004) Clin Cancer Res 10, 5708-16.
- 7. Valmori, D. et al. (2001) *Cancer Res* 61, 509-12.
- 8. Chianese-Bullock, K.A. et al. (2005) J Immunol 174, 3080-6.
- 9. Dutoit, V. et al. (2001) Cancer Res 61, 5850-6.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry

milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

3/23/24. 11:22 AM

Trademarks and Patents

Limited Uses

MAGE-A10 Antibody (#68978) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.