

#67495 Store at +4°C

## Cleaved PARP (Asp214) (D6X6X) Rabbit mAb (PE Conjugate)



**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> FC-FP	<b>Reactivity:</b> M R	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P11103	<b>Entrez-Gene Id:</b> 11545
-------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	---------------------------------

<b>Product Usage Information</b>	<b>Application</b> Flow Cytometry (Fixed/Permeabilized)	<b>Dilution</b> 1:50
<b>Storage</b>	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
<b>Specificity / Sensitivity</b>	Cleaved PARP (Asp214) (D6X6X) Rabbit mAb (PE Conjugate) recognizes endogenous levels of the large fragment (89 kDa) of rodent PARP protein only when cleaved at Asp214.	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp214 of rodent PARP1 protein.	
<b>Product Description</b>	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Cleaved PARP (Asp214) (D6X6X) Rabbit mAb #94885.	
<b>Background</b>	PARP, a 116 kDa nuclear poly (ADP-ribose) polymerase, appears to be involved in DNA repair in response to environmental stress (1). This protein can be cleaved by many ICE-like caspases <i>in vitro</i> (2,3) and is one of the main cleavage targets of caspase-3 <i>in vivo</i> (4,5). In human PARP, the cleavage occurs between Asp214 and Gly215, which separates the PARP amino-terminal DNA-binding domain (24 kDa) from the carboxy-terminal catalytic domain (89 kDa) (2,4). PARP helps cells to maintain their viability; cleavage of PARP facilitates cellular disassembly and serves as a marker of cells undergoing apoptosis (6).	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Satoh, M.S. and Lindahl, T. (1992) <i>Nature</i> 356, 356-358.</li> <li>2. Lazebnik, Y. A. et al. (1994) <i>Nature</i> 371, 346-347.</li> <li>3. Cohen, G.M. (1997) <i>Biochem. J.</i> 326, 1-16.</li> <li>4. Nicholson, D. W. et al. (1995) <i>Nature</i> 376, 37-43.</li> <li>5. Tewari, M. et al. (1995) <i>Cell</i> 81, 801-809.</li> <li>6. Oliver, F.J. et al. (1998) <i>J. Biol. Chem.</i> 273, 33533-33539.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Applications Key</b>	<b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized)
<b>Cross-Reactivity Key</b>	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.  Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner

that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.